

Quasars in the 2MASS Second Incremental Data Release

Wayne A. Barkhouse and Patrick B. Hall¹

Department of Astronomy, University of Toronto,

60 St. George Street, Toronto, ON, M5S 3H8

barkhous@astro.utoronto.ca

Received _____; accepted _____

¹Current Affiliations: Princeton University Observatory, Princeton, NJ 08544-1001 and Pontificia Universidad Católica de Chile, Departamento de Astronomía y Astrofísica, Facultad de Física, Casilla 306, Santiago 22, Chile; E-mail: phall@astro.puc.cl

ABSTRACT

Using the 2MASS Second Incremental Data Release, we have searched for near infrared counterparts to 13214 quasars from the Veron-Cetty & Veron (2000) catalog. We have detected counterparts within $4''$ for 2277 of the approximately 6320 quasars within the area covered by the 2MASS Second Incremental Data Release. Only 1.6% of these are expected to be chance coincidences. Though this sample is heterogeneous, we find that known radio-loud quasars are more likely to have large near-infrared-to-optical luminosity ratios than radio-quiet quasars are, at a statistically significant level. This is consistent with dust-reddened quasars being more common in radio-selected samples than in optically-selected samples, due to stronger selection effects against dust-reddened quasars in the latter. We also find a statistically significant dearth of optically luminous quasars with large near-infrared-to-optical luminosity ratios. This can be explained in a dust obscuration model but not in a model where synchrotron emission extends from the radio into the near-infrared and creates such large ratios. We also find that selection of quasar candidates from the $B - J/J - K$ color-color diagram, modelled on the $V - J/J - K$ selection method of Warren, Hewett & Foltz (2000), is likely to be more sensitive to dust-obscured quasars than selection using only infrared-infrared colors.

Subject headings: catalogs — infrared: general — quasars: general — galaxies: active

1. Introduction

Active Galactic Nuclei (AGN) have been studied for several decades, but our understanding of many aspects of AGN is still rudimentary. The observed emission from AGN is thought to be powered by the release of gravitational potential energy from matter falling into supermassive black holes, but the details of this process and their effect on the observed spectra of AGN are not understood. Our inability to definitively understand AGN to date may be due in part to selection effects. Most high-luminosity AGN (quasars) have been selected using their rest-frame ultraviolet through optical colors in optically magnitude-limited samples. It is therefore almost certain that dust obscuration has masked some portion of the AGN population from detection, and that AGN found by their ultraviolet-optical colors are less dusty than the truly typical AGN (Webster et al. 1995; Baker 1997; Wills & Hines 1997). This bias against dust-obscured objects can affect our understanding of AGN, in particular their connection to their host galaxies and nuclear environments. (We use the term dust *obscuration* to refer to the combined effects of dust *extinction*, the reduction in observed flux from an object screened by dust, and dust *reddening*, which arises from typical ‘non-gray’ extinction that is stronger at bluer wavelengths. Extinction can occur without reddening, but reddening cannot occur without extinction.)

The Two-Micron All-Sky Survey (2MASS, Skrutskie et al. 1997) has now made it feasible to select samples of AGN in the near infrared (NIR). A sample selected only by NIR magnitude is an excellent way to reduce any bias against dust-obscured objects, but for efficient selection of AGN candidates it is usually necessary to use additional selection criteria such as morphology and color. To aid in this endeavor and to provide a list of AGN with homogeneous infrared data, in this paper we present the results of a cross-correlation of the 2MASS Second Incremental Data Release with the quasar catalog of Veron-Cetty &

Veron (2000, hereafter VV00).

2. Catalog Cross-Correlation

We took all 13214 quasars (defined as AGN more luminous than $M_B = -23$, regardless of radio power) from Table 1 of VV00 and searched for unresolved 2MASS counterparts within $5''$. We found 2327 counterparts for 2320 quasars and discarded the more distant ones for the seven quasars with two counterparts. There were 4000 additional quasars which fell within the 2MASS areal coverage but were not detected by it.

The internal errors on the 2MASS astrometry are on average $0''.3$ for our detections, so the uncertainty in the astrometric matching is dominated by the errors on the quasar coordinates. Only 53 of the VV00 quasars have positions which may be uncertain by several arcminutes (denoted ‘A’ by VV00). The majority (8884) have optical positions which are accurate to $\lesssim 1''$ (‘O’), and 523 have similarly accurate radio positions (‘R’). The remaining 3754 have positions which should be good to a few arcsec (these have no code in VV00, but we denote them here by ‘N’).

A search for counterparts to fake objects indicates that even our near-IR counterparts of code ‘N’ objects are likely to be real. We shifted the declinations of all 13214 VV00 quasars $6'$ to the north and searched for matches within $5''$ to determine the number of spurious counterparts in our sample. Only 74 fake-object counterparts were found, as opposed to 2320 real-object counterparts. This indicates that there are few spurious counterparts in our sample, but the contamination can be reduced even further by requiring a match within $4''$ instead of $5''$. Only 43 of 2320 counterparts (including the one code ‘A’ object among the 2320) have a separation between the VV00 and 2MASS positions of $> 4''$, while 37 of 74 fake-object matches have separations $> 4''$. Thus the number of counterparts

to VV00 quasars with separations between $4''$ and $5''$ is consistent with the number expected from random chance, and for our final sample we only consider counterparts with separations $< 4''$. Our final catalog consists of 2MASS counterparts to 2277 VV00 quasars, of which about 37 (1.6%) are expected to be chance coincidences. The 2MASS and VV00 identifications of these 2277 quasars are given in Table 1. The 4000 VV00 quasars not detected by 2MASS despite lying within the area of the Second Incremental Data Release are listed in Table 2.

3. Properties of Quasars Detected by 2MASS

Radio luminosities were computed for all VV00 quasars that had measured 6 cm radio flux densities. We assumed a radio spectral index of $\alpha = -0.5$, a Hubble constant of $H_o = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $q_o = 0.0$. We classified each quasar as radio-loud (RLQ) or radio-quiet (RQQ) based on the ratio of its radio to optical luminosity (L_{rad}/L_{opt}). The divide between RLQ and RQQ was taken as $\log(L_{rad}/L_{opt})=0$. Optical luminosities were calculated from M_B as given in VV00. Near infrared luminosities were calculated from K magnitudes given in the 2MASS catalog using k -corrections for a power-law continuum with $\alpha = -0.5$, the same value used by VV00 to calculate their M_B values.

Figure 1 shows the degeneracy between luminosity and redshift inherent in any single flux-limited survey. However, it is clear that 2MASS can in principle detect quasars to at least $z = 5$.

Figure 2 shows that most VV00 quasars with radio information are radio-loud quasars, since VV00 do not report upper limits on radio flux densities. Assuming that VV00 reported radio data for all radio-selected quasars, the majority of radio-selected quasars catalogued in VV00 are radio-loud, and most radio-quiet quasars in VV00 must not have

been radio-selected. The RLQ vs. RQQ bimodality seen in heterogeneous quasar samples is not erased by plotting L_{rad}/L_{NIR} instead of L_{rad}/L_{opt} . However, it is interesting to note that some objects which qualify as radio-loud using L_{rad}/L_{opt} would be classified as radio-quiet using L_{rad}/L_{NIR} (or vice-versa, depending on where the division between RLQ and RQQ is made in L_{rad}/L_{NIR}). One possible explanation for this might be because some radio-selected quasars in VV00 are obscured by dust and have low values of L_{opt} for their L_{NIR} .

Figure 3 shows that radio-loud quasars from VV00 are more likely to have large L_{NIR}/L_{opt} ratios ($L_{NIR}/L_{opt} \gtrsim 1$) than radio-quiet quasars from VV00. 9.9% of known RLQs from VV00 which are detected in 2MASS have $L_{NIR}/L_{opt} \geq 1$ compared to only 2.5% of RQQs and quasars with unknown radio properties. Again, this is understandable if some radio-selected quasars in VV00 are obscured by dust and consequently have their values of L_{opt} suppressed relative to their L_{NIR} . Radio-quiet quasars similarly obscured by dust would be underrepresented in magnitude-limited optically selected samples, especially color-selected ones, and thus less likely to be catalogued in VV00. Note that the many upper limits at large L_{NIR}/L_{opt} belong to optically faint quasars whose L_{NIR} values are poorly constrained by the relatively shallow 2MASS data. Also note the shifts toward slightly larger L_{NIR}/L_{opt} at $z < 0.5$ and $2.1 \lesssim z \lesssim 2.5$. The former is probably due to host galaxy contributions to L_{NIR} , and the latter to the presence of $H\alpha$ in the K_s band, neither of which are compensated for by our simple power-law k -correction.

To determine if the L_{NIR}/L_{opt} distribution of the RLQs differed significantly from that of the other quasars, i.e. the RQQs and quasars with unknown radio properties (RUQs), we used the Peto-Prentice Generalized Wilcoxon test for censored datasets (Feigelson & Nelson 1985), as implemented in the STSDAS.STATISTICS package of IRAF.² This test is

²The Image Reduction and Analysis Facility (IRAF) is distributed by the National Optical

insensitive to differences in the censoring patterns for the two samples being compared. The combination of RQQs and RUQs forms a valid, conservative sample for comparison with the RLQs, since RQQs comprise $\sim 90\%$ of quasars and since any unrecognized RLQs among the RUQs will dilute any intrinsic difference between the L_{NIR}/L_{opt} distributions of RLQs and RQQs. We find that the L_{NIR}/L_{opt} ratios distributions of these two samples are different at the 99.99% confidence level. Again, we do not believe that the higher fraction of large L_{NIR}/L_{opt} ratio objects among *known* RLQs compared to *known* RQQs is an intrinsic property of RLQs, but is merely a reflection of the lesser bias against dust-obscured quasars with large L_{NIR}/L_{opt} ratios in radio-selected samples compared to optically-selected samples.

Figure 4 corroborates the suggestion that quasars with large L_{NIR}/L_{opt} ratios are obscured by dust. It shows that such quasars are much less common among the population of quasars more luminous than $M_B \simeq -27$ than among the less luminous population. The Peto-Prentice test shows that this difference is significant at $\geq 99.995\%$ confidence whether the sample of all VV00 quasars is split at $M_B = -27$ or is split in half (at $M_B = -26.2$). This is to be expected if dust obscuration is lowering L_{opt} (and M_B) relative to L_{NIR} , thus increasing L_{NIR}/L_{opt} . By itself this does not rule out alternative explanations. However, host galaxy contributions to the NIR luminosities would require extremely luminous host galaxies since the quasars with large L_{NIR}/L_{opt} ratios have $0.5 < z < 3$ (Figure 3). Also, a scenario where large L_{NIR}/L_{opt} ratios are caused by synchrotron emission extending into the NIR in some radio-loud quasars (Serjeant & Rawlings 1997) does not explain why such emission should preferentially occur in less luminous quasars. Dust obscuration seems the simplest explanation for Figure 4, although we note that Francis, Whiting & Webster

(2000) concluded that synchrotron emission was the best explanation for the red colors of the many RLQs in the Parkes Half-Jansky sample of Webster et al. (1995).

4. Selection of Quasars Using 2MASS

As discussed in the introduction, if there indeed exists a population of dust-obscured radio-loud quasars, then it is desirable to search for a population of dust-obscured radio-quiet quasars without being strongly biased against such objects. Selection on NIR magnitudes reduces bias due to dust extinction, and selection on NIR colors reduces bias due to dust reddening. We use the sample of VV00 quasars detected by 2MASS to investigate NIR selection. However, it must be kept in mind that the majority of quasars from VV00 are probably not strongly dust-obscured due to their predominantly optical selection. The range of quasar colors in our large sample may reflect the true range of quasar colors, but the distribution of colors will not reflect the true distribution.

4.1. Selection on Near-Infrared Colors

Nelson et al. (1998) have begun identifying candidate obscured AGN in 2MASS using a simple criterion of $J - K \geq 2$, as reported in Cutri et al. (2000) and Smith et al. (2000b). This criterion was chosen based on the colors of a sample of miscellaneous AGN including the Palomar-Green (PG) AGN (Schmidt & Green 1983; Neugebauer et al. 1987), from which Cutri et al. (2000) conclude that “virtually all of these known QSOs have $J - K \leq 2$” However, as seen in Figure 5, a $J - K \geq 2$ criterion selects 13% of quasars from VV00 at $z \leq 0.5$, as well as a smaller percentage at $0.5 < z < 3$. Thus, unless the known quasars with $J - K \geq 2$ in VV00 are preferentially dust-obscured quasars, a $J - K \geq 2$ selection criterion will suffer some contamination from unobscured quasars. On the other hand, all

but four of the 2277 VV00 quasars detected by 2MASS have $J - K \leq 2.5$, whereas the 2MASS AGN of Cutri et al. (2000) have values as red as $J - K = 3.5$. Therefore a selection criterion of $J - K \geq 2.5$ should select dust-obscured AGN with little contamination.

However, a $J - K \geq 2.5$ selection criterion would not be complete; that is, it would not sample the full population of obscured AGN. We know this for two reasons. First, Cutri et al. (2000) do not find evidence of dust reddening only in AGN with $J - K \geq 2.5$. Second, we show in the next section that $J - K$ color is not well correlated with $B - K$ color, whose longer wavelength baseline than any optical-NIR color besides $U - K$ makes it arguably the most efficient photometric measure of the reddening of a quasar in the optical-NIR (Webster et al. 1995), given the difficulty in obtaining deep wide-field U -band data. A long wavelength baseline is needed to distinguish reddening from the intrinsic dispersion in quasar colors. The observed dispersion in the 2MASS quasar sample is only $0^{\text{m}}31$ in $J - K$, vs. $0^{\text{m}}67$ in $B - K$ (calculated using 5σ clipping to remove outliers, and using only $1 < z < 2$ quasars for $J - K$). However, the sensitivity to reddening is given by the ratio of the color excess to this dispersion: $E(J - K)/\sigma_{J-K}=2.12$ vs. $E(B - K)/\sigma_{B-K}=3.84$ at $z = 3$ (the difference is even more extreme at $z = 0$). Simply put, less extinction is required to produce anomalously red colors in $B - K$ than in $J - K$, or indeed than in any other optical-NIR color besides $U - K$. Finally, the variation of $J - K$ color with redshift seen in Figure 5 means that a simple $J - K$ color cut will not select reddened AGN with equal efficiency at all redshifts. Thus the evidence available to date, while scarce, suggests that infrared colors alone are insufficient to *efficiently* sample the full parameter space open to obscured quasars.

4.2. Selection on Optical and Near-Infrared Colors

The 2MASS catalog includes B and R magnitudes from the USNO-A and Tycho catalogs, as well as JHK . As noted by Francis et al. (2000) and Cutri et al. (2000), sampling the full parameter space open to obscured quasars will be difficult based on $B - R$ vs. $J - K$ colors. However, Warren et al. (2000, hereafter WHF) have proposed a promising criterion for selecting K -band flux-limited samples of quasars from the $V - J$ vs. $J - K$ color-color diagram, which they dub the KX method. We do not have V magnitudes for the majority of our quasars, but the $B - J$ vs. $J - K$ diagram (Figure 6) shows the same gross features as the $V - J$ vs. $J - K$ diagram of WHF (their Figure 2). Note that we plot our diagram upside-down relative to theirs, however. The dotted points in Figure 6 are an essentially random sample of 14,000 high Galactic latitude objects. The filled squares are the 2277 VV00 quasars with 2MASS counterparts within $4''$: known radio-quiet quasars among them are denoted by overplotted boxes, and known radio-loud quasars by overplotted crosses. The bulk of known quasars have blue $B - J$ colors, but there is a tail of candidate obscured quasars, predominantly RLQs, with $J - K \sim 1.75$ and $B - J \geq 2$. Note that most of these quasars do not have $J - K > 2$. The dashed lines of constant $B - K$ color show that red $J - K$ color does not correlate strongly with red $B - K$ color either.

We cannot rule out the possibility that $J - K$ does correlate with $B - J$ or $B - K$ but that this correlation is masked by strong selection effects against quasars with red $B - J$ and $J - K$ in Figure 6. However, we have also plotted a subsample of radio-selected quasars from the completely identified Parkes Half-Jansky sample (triangles; Francis et al. 2000). There is no strong correlation between $J - K$ and $B - J$ or $B - K$ for these objects either. Selecting objects red in $J - K$ would not appear to be an efficient way of selecting objects red in $B - K$. With RLQs, there is no guarantee that such red $B - K$ colors are due to dust rather than synchrotron emission with a short wavelength cutoff in the rest-frame

optical (Serjeant & Rawlings 1997; Francis et al. 2000). However, with RQQs the latter effect will be much weaker and any confirmed population of RQQs with $B - K \gtrsim 4$ will almost certainly be reddened by dust.

The $R - J$ vs. $J - K$ color-color diagram (Figure 7) is similar to the $V - J$ and $B - J$ vs. $J - K$ diagrams, and as pointed out by WHF offers the advantage of extending the effectiveness of the KX method from $z \simeq 3.4$ to $z \simeq 4.4$. Beyond $z \simeq 4.4$, $\text{Ly}\alpha$ forest absorption greatly depresses the flux in the R band, and the resulting red $R - J$ colors may result in confusion with stars.

We have constructed selection criteria similar to the $V - J/J - K$ criterion of WHF to illustrate the separation of quasars from stars in both $B - J/J - K$ and $R - J/J - K$ color-color diagrams (solid lines in Figure 6 and Figure 7). We adopt the following selection criteria in $B - J/J - K$:

$$(J - K) \geq 0.3 + 0.2 \times (B - J) \quad (1)$$

and the following in $R - J/J - K$:

$$(J - K) \geq 0.4 + 0.4 \times (R - J) \quad (\text{for } R - J < 1.625) \quad (2)$$

$$(J - K) \geq 1.05 \quad (\text{for } R - J \geq 1.625) \quad (3)$$

As pointed out by WHF and as shown by the arrows in Figure 6 and Figure 7, reddening moves quasars approximately parallel to these selection criteria and the bluer portion of the stellar locus in these diagrams, resulting in good sensitivity to reddened and unreddened quasars. Only 124 of 2277 quasar counterparts (5.4%) have colors that do not meet the $B - J/J - K$ selection criteria, while 283 (12.4%) have colors that do not meet the $R - J/J - K$ selection criteria. Some of these are probably chance coincidences of stars with VV00 quasars. Assuming all 37 estimated spurious matches are among them, then only 87 quasar counterparts (3.8%) do not meet the $B - J/J - K$ selection criteria while

236 quasar counterparts (10.4%) do not meet the $R - J/J - K$ selection criteria. (Since these criteria are illustrative only, we do not consider how efficient they are at rejecting stars, but any application of them to 2MASS or other data must consider how close to the stellar locus the selection criteria can be drawn as a function of magnitude without selecting too many spurious candidates.)

For both selection criteria, the selection efficiency is dependent on redshift. Figure 8a shows our $B - J/J - K$ selection parameter versus z for all VV00 quasar counterparts, while Figure 8b shows our $R - J/J - K$ selection parameter. The selection parameter is essentially the distance in the color-color diagram to the selection criterion line which divides quasar candidates from stars. Objects with a positive selection parameter are accepted as candidates from that diagram. While $R - J/J - K$ selection may be more useful at $z \geq 3.4$ than $B - J/J - K$ selection due to Ly α forest absorption in B , it is less efficient at separating quasars from stars at $z \leq 3.4$. However, our $B - J/J - K$ criterion is not perfect: the intrinsic dispersion in quasar colors plus the redshift dependence of quasar colors may result in reduced selection efficiency at $1 < z < 2$ compared to $z < 0.5$ and $2 < z < 2.5$.

Thus, the $V - J/J - K$ criterion of WHF, or a $B - J/J - K$ selection criterion patterned after theirs, appear to be a more complete method of selecting reddened AGN than the simple $J - K > 2$ selection of Nelson et al. (1998). Nevertheless, Smith et al. (2000a) find that the 2MASS AGN of Nelson et al. (1998) span a much larger range in $B - K$ color and in optical polarization than the Palomar-Green (PG) AGN even when the comparison is restricted to PG AGN with $J - K > 2$. Both of these trends are consistent with a much higher frequency of dust-obscured AGN in the 2MASS $J - K > 2$ selected sample. This high frequency of dust-obscured AGN found using a simple and probably incomplete criterion is unexpected given the lack of a correlation between $J - K$ and $B - J$

(or $B - K$), but it does suggest that more complete $B - J/J - K$ selected AGN samples will be very efficient at finding great numbers of dust-obscured AGN. Of course, a key test will be to see where the 2MASS AGN of Nelson et al. (1998) fall in the $B - J/J - K$ diagram.

5. Conclusions

We have identified near infrared counterparts within $4''$ of 2277 quasars from the Veron-Cetty & Veron (2000) catalog using the 2MASS Second Incremental Data Release. Approximately 6320 quasars from Veron-Cetty & Veron (2000) were located within the area covered by the 2MASS catalog. This detection rate of 36% suggests that the final 2MASS catalog will yield matches for approximately 4760 of the 13214 VV00 quasars. Our sample of 2277 quasars with IR data is nearly ten times larger than the largest previous such sample (Srianand & Kembhavi 1997), and more than ten times larger than the largest previous samples with homogeneous IR data (Warren et al. 2000; Francis et al. 2000).

The observation that the fraction of radio-loud quasars with large L_{NIR}/L_{opt} ratios is higher than the fraction of probable radio-quiet quasars with such ratios supports the suggestion that *known* radio-selected quasars are more likely to be obscured by dust than *known* optically-selected quasars due to stronger selection effects against dust-obscured quasars in optically selected samples. Known quasars with large L_{NIR}/L_{opt} ratios are also underrepresented among optically luminous quasars ($M_B < -27$). This is again consistent with a selection effect caused by dust obscuration, which would tend to lower L_{opt} and M_B while increasing L_{NIR}/L_{opt} . These results need to be confirmed using homogeneous quasar samples rather than the heterogeneous VV00 sample, but the statistical significance of our results in such a heterogeneous sample bodes well for their reality in more carefully selected samples.

Following Warren, Hewett & Foltz (2000), we show that the $B - J/J - K$ color-color diagram is more sensitive to the detection of dust-obscured quasars than using infrared colors alone. The stage is now set for the determination of the true fraction of dust-obscured RQQs using large NIR surveys such as 2MASS and DENIS. The detection of dust-obscured AGN will also benefit greatly from hard X-ray surveys, but for the foreseeable future NIR selection will be possible over much larger areas than hard X-ray selection, and thus more sensitive to luminous AGN which are more easily studied in detail.

We thank Željko Ivezić and the referee Paul Francis for useful comments, and Roc Cutri for determining which quasars were observed but not detected by 2MASS. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

REFERENCES

- Allen, C. W. 2000, *Astrophysical Quantities* (New York: Springer-Verlag), 527
- Baker, J. 1997, *MNRAS*, 286, 23
- Cutri, R. M., Nelson, B. O., Kirkpatrick, J. D., Huchra, J. P. & Smith, P. S. 2000, to appear in *The New Era of Wide-Field Astronomy*, ed. R. Clowes (San Francisco: ASP)
- Feigelson, E. D., & Nelson, P. I. 1985, *ApJ*, 293, 192
- Francis, P., Whiting, M. & Webster, R. 2000, *PASA*, 17, 56
- Nelson, B., Cutri, R. M., Kirkpatrick, J. D., Beichman, C. A., Lonsdale, C., Huchra, J. P. & Skrutskie, M. F. 1998, in *American Astronomical Society Meeting*, Vol. 193, 8104
- Neugebauer, G., Green, R. F., Matthews, K., Schmidt, M., Soifer, B. T. & Bennett, J. 1987, *ApJS*, 63, 615
- Schmidt, M. & Green, R. F. 1983, *ApJ*, 269, 352
- Serjeant, S. & Rawlings, S. 1997, *Nature*, 379, 304
- Skrutskie, M. F., Schneider, S. E., Stiening, R., Strom, S. E., Weinberg, M. D., Beichman, C., Chester, T., Cutri, R., Lonsdale, C., Elias, J., Elston, R., Capps, R., Carpenter, J., Huchra, J., Liebert, J., Monet, D., Price, S. & Seitzer, P. 1997, in *The Impact of Large Scale Near-IR Sky Surveys*, eds. F. Garzon et al. (Dordrecht: Kluwer), 25
- Smith, P. S., Schmidt, G. D., Hines, D. C., Cutri, R. M. & Nelson, B. O. 2000a, in *American Astronomical Society Meeting*, Vol. 196, 5014
- Smith, P. S., Schmidt, G. D., Hines, D. C., Cutri, R. M. & Nelson, B. O. 2000b, *ApJ*, in press (astro-ph/0010173)

- Srianand, R. & Kembhavi, A. 1997, *ApJ*, 478, 70
- Veron-Cetty, M. P. & Veron, P. 2000, *European Southern Observatory Scientific Report*, 19, 1
- Warren, S. J., Hewett, P. C. & Foltz, C. B. 2000, *MNRAS*, 312, 827
- Webster, R. L., Francis, P. J., Peterson, B. A., Drinkwater, M. J. & Masci, F. J. 1995, *Nature*, 375, 469
- Wills, B. J. & Hines, D. C. 1997, in *Mass Ejection from Active Galactic Nuclei*, eds. N. Arav, I. Shlosman, and R. J. Weymann (ASP: San Francisco), 99




Fig. 1.— This figure shows the correlation between the NIR luminosity and the redshift of all quasars in VV00 that fall within the area measured by 2MASS. RLQs (crosses) and RQQs (filled squares) are plotted as well as quasars which did not have measured radio flux density in VV00 (“Radio Unknown”; circled dots). We have also included quasars which did not have a corresponding detection in 2MASS (“2MASS upper Limit”; dots) and have used a typical high Galactic latitude 2MASS limit of $K = 15.50$ to compute the NIR luminosity.




Fig. 2.— This figure shows the relationship between the ratio of $L_{\text{rad}}/L_{\text{NIR}}$ as a function of redshift for RLQs (crosses) and RQQs (filled squares). Most of the quasars in this plot are radio-loud because most RQQs have only upper limits to their radio fluxes, which are not reported by VV00.




Fig. 3.— This figure illustrates the relationship between the ratio of $L_{\text{NIR}}/L_{\text{opt}}$ versus redshift for the sample of quasars in VV00 which lie within the 2MASS coverage area. Symbols are the same as in Figure 1. Radio-loud quasars are more likely to have large $L_{\text{NIR}}/L_{\text{opt}}$ ratios than radio-quiet or radio-unknown quasars. The peak value of the $L_{\text{NIR}}/L_{\text{opt}}$ distribution is shifted slightly higher at $z < 0.5$ probably because of host galaxy contributions to L_{NIR} , and at $z \sim 2.3$ probably because of $H\alpha$ contributions to L_{NIR} . Neither of these effects are compensated for by our simple power-law k -correction. The handful of $z > 3.5$ objects suggest a shift toward larger $L_{\text{NIR}}/L_{\text{opt}}$ at those redshifts as well, which could be due to a systematic inaccuracy below the Lyman limit in the k -correction applied by VV00 in their calculation of M_B . Such a shift would not affect our conclusions, however, given the small number of objects in VV00 at $z > 3.5$. The many upper limits at large $L_{\text{NIR}}/L_{\text{opt}}$ belong to optically faint quasars whose L_{NIR} values are poorly constrained by the relatively shallow 2MASS data.




Fig. 4.— The ratio of the near-infrared to optical luminosity is shown as a function of absolute B magnitude for the sample of quasars in VV00 which lie within the 2MASS coverage area. Symbols are the same as in Figure 1. The M_B values from VV00 are accurate only to ± 0.1 , so a random offset between -0.04 and $+0.05$ magnitudes has been added to them to spread out the points in the figure. The peak value of the $L_{\text{NIR}}/L_{\text{opt}}$ distribution is shifted slightly higher at low luminosities, probably because of host galaxy contributions to L_{NIR} in the low redshift objects which dominate the sample at low luminosities. The fact that most quasars with large $L_{\text{NIR}}/L_{\text{opt}}$ ratios are fainter than $M_B \simeq -27$ is consistent with the suggestion that quasars with large $L_{\text{NIR}}/L_{\text{opt}}$ ratios are obscured by dust. The many upper limits at large $L_{\text{NIR}}/L_{\text{opt}}$ belong to optically faint quasars whose L_{NIR} values are poorly constrained by the relatively shallow 2MASS data.




Fig. 5.— $J - K$ versus redshift for all 2MASS counterparts of VV00 quasars (small filled squares). Known radio-loud quasars are denoted by overplotted crosses. Known radio-quiet quasars are denoted by overplotted boxes; they are found only at low redshift because VV00 do not list upper limits to radio fluxes, and thus more distant radio-quiet quasars form the bulk of the “radio unknown” population. The solid line shows the $J - K > 2$ selection criterion of Nelson et al. (1998).




Fig. 6.— $B - J$ vs. $J - K$ color-color diagram for all 2MASS counterparts of VV00 quasars. The small dots are approximately 14,000 essentially random $K \leq 15$ 2MASS point sources. They illustrate the colors of the stellar locus as well as other objects (quasars and compact galaxies) seen at moderate to high Galactic latitude. The small filled squares are the 2277 VV00 quasars with 2MASS counterparts within $4''$. Known radio-quiet quasars are denoted by overplotted boxes, and known radio-loud quasars are denoted by overplotted crosses. Dashed lines show lines of constant $B - K$ color. Radio-loud quasars outnumber radio-quiet or radio-unknown quasars for $B - K \gtrsim 3.5$, whereas the opposite is true for $B - K \lesssim 3.5$. This heterogeneous sample of RLQs from VV00 has a range of colors similar to the well-defined Parkes sample of Francis, Whiting & Webster (2000; open triangles). The two arrows show the reddening vectors for one magnitude of visual extinction in the quasar rest frame ($A_V = 1$, or $E(B - V) = 0.32$; Table 21.6 of Allen 2000). The shorter arrow is for $z = 0$ and the longer arrow is for $z = 3$. The solid line shows the selection criteria we adopt to separate stars from quasar candidates; note that reddening moves quasars approximately parallel to the selection criteria.




Fig. 7.— $R - J$ vs. $J - K$ color-color diagram for all 2MASS counterparts of VV00 quasars. The small dots are approximately 14,000 essentially random $K \leq 15$ 2MASS point sources. The small filled squares are the 2277 VV00 quasars with 2MASS counterparts within $4''$. The open triangles are the Parkes quasars of Francis, Whiting & Webster (2000). The two arrows show the reddening vectors for one magnitude of visual extinction in the quasar rest frame ($A_V = 1$, or $E(B - V) = 0.32$; Table 21.6 of Allen 2000). The shorter arrow is for $z = 0$ and the longer arrow is for $z = 3$. The solid line shows the selection criteria we adopt to separate stars from quasar candidates; note that reddening moves quasars approximately parallel to the selection criteria.




Fig. 8.— Quasar selection criteria parameters versus redshift for all 2MASS counterparts of VV00 quasars. In a) we show the $B-J/J-K$ selection parameter $(J-K)-0.3-0.2\times(B-J)$, and in b) we show the analogous parameter for $R-J/J-K$ selection. In both cases this parameter is the shortest distance in color-color space to the selection criterion line which divides quasar candidates from stars. Objects with positive values of these quantities (above the solid lines) would be accepted as quasar candidates. The intrinsic dispersion and redshift dependence of quasar colors means that the selection efficiency will be redshift dependent, but overall $B-J/J-K$ selection is more efficient than $R-J/J-K$ selection.

Table 1. VV00 Quasars Detected in 2MASS^a

2MASS ID	VV00 ID	RA deg	DEC deg	L_{NIR} Jy	L_{opt} Jy	b-m mag	r-m mag	J mag	σ_J mag	H mag	σ_H mag	K mag	σ_K mag	M_B mag	dist _{opt} arcsec	Radio Jy	z
0000029-350332	MS 23574-3520	0.012084	-35.059052	24.14	23.91	17.20	17.10	16.221	0.110	15.332	0.119	14.513	0.099	-25.50	0.14	-99.00	0.508
0000244-124548	PHL 2525	0.101767	-12.763336	24.06	23.71	15.10	14.80	14.475	0.034	13.704	0.041	12.580	0.034	-25.00	0.31	-99.00	0.200
0001323+211336	TEX 2358+209	0.384921	21.226782	24.89	23.87	18.20	18.60	16.177	0.102	15.547	0.142	14.526	0.093	-25.40	0.25	26.93	1.106
0002224-000444	PB 5694	0.593673	-0.078939	24.08	23.39	18.60	18.30	16.586	0.122	16.500	0.259	15.775	0.229	-24.20	0.79	-99.00	0.810
0003153-275157	HE 0000-2808	0.813904	-27.865856	24.68	24.55	17.60	16.50	15.996	0.091	15.653	0.160	14.944	0.131	-27.10	0.43	-99.00	1.051

^aThe complete version of this table is in the electronic edition of the Journal or at www.astro.utoronto.ca/~barkhous/quasar_table1.gz. The printed edition contains only a sample.

Note. — VV00 IDs are the Name entries of VV00. A handful of objects which have blank Name entries in VV00 are identified here by the prefix VV followed by their coordinates in VV00. Objects having two 2MASS counterparts within 4'' are identified by an asterisk preceding their 2MASS ID number. Only the closest counterpart is listed here. Columns 3 and 4 give the coordinates for J2000. Columns 5 and 6 lists the logarithm of the calculated NIR and optical luminosity. Column 16 gives the distance to the associated optical source. Column 17 lists the radio flux density measured at 6 cm. Data for columns 2, 15, 17, and 18 are from VV00, while the remaining data are from 2MASS. Unavailable data entries are given as -99.00.

Table 2. VV00 Quasars With Only Upper Limits in 2MASS^a

VV00 ID	RA	DEC	L_{opt}	Radio	z
	deg	deg	Jy	Jy	
PKS 2357-326	0.084167	-32.350278	24.03	27.53	1.275
PKS 2358-161	0.272500	-15.852222	24.67	28.01	2.044
TEX 2358+189	0.285833	19.242778	24.63	28.30	3.100
UM 195	0.442500	-1.992778	23.71	-99.00	0.867
UM 196	0.458333	-1.994444	24.91	-99.00	2.810

^aThe complete version of this table is in the electronic edition of the Journal or can be found at www.astro.utoronto.ca/~barkhous/quasar_table2.gz. The printed edition contains only a sample.

Note. — IDs are the Name entries of VV00. A handful of objects which have blank Name entries in VV00 are identified here by their coordinates and the prefix VV.

Columns 2 and 3 give the coordinates for J2000. Columns 4 lists the logarithm of the calculated optical luminosity. Column 5 gives the radio flux density measured at 6 cm.

Unavailable data entries are given as -99.00.